ELSEVIER

Environmental Pollution 137 (2005) I–V

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 137

Number 1 2005

1 **Preface**
R.W. Battarbee, A. Jenkins

3 **The United Kingdom Acid Waters Monitoring Network: A review of the first 15 years and introduction to the special issue**
D.T. Monteith, C.D. Evans

After 15 years of the UK Acid Waters Monitoring Network, we can now draw clear conclusions regarding the impact of emission reductions on acidified UK fresh waters.

15 **Changes in the atmospheric deposition of acidifying compounds in the UK between 1986 and 2001**
D. Fowler, R.I. Smith, J.B.A. Muller, G. Hayman, K.J. Vincent

Deposition of acidity and sulphur in the UK has halved during the last 16 years, and substantial non-linearities in emission-deposition relationships have emerged.

27 **Trends in surface water chemistry of acidified UK Freshwaters, 1988–2002**
J.J.L. Davies, A. Jenkins, D.T. Monteith, C.D. Evans, D.M. Cooper

Acidified UK freshwaters show recovery but there are key uncertainties with regard to DOC and ANC.

41 **Evidence of sulphur and nitrogen deposition signals at the United Kingdom Acid Waters Monitoring Network sites**
D.M. Cooper

There is strong evidence of a direct link between reductions in atmospheric deposition of sulphur and reductions in sulphate concentrations in UK freshwaters at an annual time scale.

55 **Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts**
C.D. Evans, D.T. Monteith, D.M. Cooper

Large increases in dissolved organic carbon concentrations across the UK may be a combined response to climate change and decreasing acid deposition.

73 **Nitrate leaching as a confounding factor in chemical recovery from acidification in UK upland waters**
C.J. Curtis, C.D. Evans, R.C. Helliwell, D.T. Monteith

With declining excess sulphate, nitrate will become the dominant agent of continued anthropogenic acidification in many UK upland waters within a decade.

83 **Biological responses to the chemical recovery of acidified fresh waters in the UK**
D.T. Monteith, A.G. Hildrew, R.J. Flower, P.J. Raven, W.R.B. Beaumont, P. Collen, A.M. Kreiser, E.M. Shilland, J.H. Winterbottom

Recently observed changes in the species composition of UK lakes and streams are consistent with chemical recovery from acidification.

103 **The ecology of acidification and recovery: Changes in herbivore-algal food web linkages across a stream pH gradient**
M.E. Ledger, A.G. Hildrew

Generalist invertebrates maintain algae-herbivore interactions in acid streams.

119 **Defining reference conditions for acidified waters using a modern analogue approach**
G.L. Simpson, E.M. Shilland, J.M. Winterbottom, J. Keay

The use of a palaeolimnological technique to identify modern ecological reference analogues for acidified lakes is demonstrated.

135 **Reconstructing pre-acidification pH for an acidified Scottish loch: A comparison of palaeolimnological and modelling approaches**
R.W. Battarbee, D.T. Monteith, S. Juggins, C.D. Evans, A. Jenkins, G.L. Simpson

Methods of reconstructing pre-acidification pH for an acidified Scottish loch compared.

151 **Temporal trends in spheroidal carbonaceous particle deposition derived from annual sediment traps and lake sediment cores and their relationship with non-marine sulphate**
N.L. Rose, D.T. Monteith

Temporal trends in spheroidal carbonaceous particles from sediment records show good agreement with non-marine sulphate deposition over various time-scales.

S0269–7491(05)00306-4

165 Regional scale evidence for improvements in surface water chemistry 1990–2001
B.L. Skjelkvåle, J.L. Stoddard, D.S. Jeffries, K. Tørseth, T. Høgåsen, J. Bowman, J. Mannio, D.T. Monteith, R. Mosello, M. Rogora, D. Rzychon, J. Vesely, WJ. Wieting, A. Wilander, A. Worsztynowicz

Data suggest that continued improvement of acid-sensitive lakes and streams will lead to biological recovery.

Publisher's note

177 Online submission and review for Environmental Pollution

Number 2 2005

New initiative

179 Mercury loss from soils following conversion from forest to pasture in Rondônia, Western Amazon, Brazil
M.D. Almeida, L.D. Lacerda, W.R. Bastos, J.C. Herrmann

Deforestation can be responsible for maintaining high Hg levels in the Amazon environment, through a grasshopper effect of Hg remobilization from the affected soils.

187 Assessment of contaminant lability during phytoremediation of polycyclic aromatic hydrocarbon impacted soil
Z.D. Parrish, M.K. Banks, A.P. Schwab

The lability of sorbed contaminants is modified by the presence of plants. Remediation coupled with plant treatment can change the bioavailability of contaminants in soil.

198 Organotin contamination in fishes with different living patterns and its implications for human health risk in Taiwan
C.-C. Lee, T. Wang, C.-Y. Hsieh, C.-J. Tien

Spatial and seasonal variations in accumulation of organotins were showed by fishes with different living patterns and the potential health risk to ingest such fishes.

209 Selenite reduction in Boom clay: Effect of FeS_2, clay minerals and dissolved organic matter
C. Bruggeman, A. Maes, J. Vancluysen, P. Vandemussele

Selenite reduction by FeS_2 is kinetically controlled, with clay minerals and organic matter playing an important role.

222 Multi-species interactions impact the accumulation of weathered 2,2-bis (*p*-chlorophenyl)-1,1-dichloroethylene (*p,p′*-DDE) from soil
J.W. Kelsey, J.C. White

Interactions between earthworms and plants affect both the phytoextraction and bioaccumulation of p,p′-DDE in soil.

231 A comparison of the rhizosphere-based method with DTPA, EDTA, $CaCl_2$, and $NaNO_3$ extraction methods for prediction of bioavailability of metals in soil to barley
M.-H. Feng, X.-Q. Shan, S. Zhang, B. Wen

The rhizosphere-based method is a preferred approach for evaluating bioavailability of metals in soil.

241 Reflectance properties and physiological responses of *Salicornia virginica* to heavy metal and petroleum contamination
P.H. Rosso, J.C. Pushnik, M. Lay, S.L. Ustin

Spectral characteristics of S. virginica were significantly correlated with symptoms of cadmium and lightweight petroleum contamination, and had a distinct signature for both types of contaminants.

253 Monitoring and toxicity of sulfonated derivatives of benzene and naphthalene in municipal sewage treatment plants
M.C. Alonso, L. Tirapu, A. Ginebreda, D. Barceló

Sulphonated organic pollutants enter sewage treatment plants and although they partly discharged into the river waters no toxic effects are observed when biological treatment is used.

263 Impacts of lawn-care pesticides on aquatic ecosystems in relation to property value
J.P. Overmyer, R. Noblet, K.L. Armbrust

The macroinvertebrate communities of suburban streams may be influenced by the toxicity of the pesticides present in the water and sediment as well as select water quality parameters.

273 The analysis of soil cores polluted with certain metals using the Box–Cox transformation
M. Meloun, M. Sáňka, P. Němec, S. Křítková, K. Kupka

A new procedure of statistical analysis, with exploratory data diagnostics and Box–Cox transformation was used.

281 Temporal variation of trace metal geochemistry in floodplain lake sediment subject to dynamic hydrological conditions
C. van Griethuysen, M. Luitwieler, J. Joziasse, A.A. Koelmans

There is strong seasonal variation in degree of metal mobilization from lake sediments.

295 Acid neutralization within limestone sand reactors receiving coal mine drainage
B.J. Watten, P.L. Sibrell, M.F. Schwartz

A new treatment method for acid mine drainage is described.

305 Sampling technologies and air pollution control devices for gaseous and particulate arsenic: A review
L. Helsen

Submicron arsenic fumes are difficult to control in conventional air pollution control devices.

316 Soil characteristics, heavy metal availability and vegetation recovery at a former metallurgical landfill: Implications in risk assessment and site restoration
E. Remon, J.-L. Bouchardon, B. Cornier, B. Guy, J.-C. Leclerc, O. Faure

Heavy metals at a former metallurgical landfill are not leachable and poorly phytoavailable.

324 Perfluorooctane sulfonic acid and organohalogen pollutants in liver of three freshwater fish species in Flanders (Belgium): Relationships with biochemical and organismal effects
P.T. Hoff, K. Van Campenhout, K. Van de Vijver, A. Covaci, L. Bervoets, L. Moens, G. Huyskens, G. Goemans, C. Belpaire, R. Blust, W. De Coen

Hepatic perfluorooctane sulfonic acid contamination in Flanders (Belgium) might affect serological endpoints in feral carp and eel.

334 Comparison of $PM_{2.5}$ carbon measurement methods in Hong Kong, China
J.C. Chow, J.G. Watson, P.K.K. Louie, L.-W.A. Chen, D. Sin

Concentrations of elemental carbon in Hong Kong are higher than in US cities, probably because of the widespread use of diesel engines.

345 Structural and functional diversity of microbial communities from a lake sediment contaminated with trenbolone, an endocrine-disrupting chemical
V. Radl, K. Pritsch, J.C. Munch, M. Schloter

The steroid hormone trenbolone affects microbial community function in a lake sediment.

354 Availability of heavy metals for uptake by *Salix viminalis* on a moderately contaminated dredged sediment disposal site
E. Meers, S. Lamsal, P. Vervaeke, M. Hopgood, N. Lust, F.M.G. Tack

Combination of energy cropping and phytoremediation with Salix sp. allows for economic revalorization of contaminated sites during the remediation process.

365 Order-of-magnitude increase of Hg in Norwegian peat profiles since the outset of industrial activity in Europe
E. Steinnes, T.E. Sjøbakk

Results from analyses of ombrotrophic peat cores support the view that the present level of Hg0 in the atmosphere is mainly of anthropogenic origin.

Number 3 2005

Editorial

371 Introductory remarks to the Special Issue
S. Huttunen

Papers

372 UV-B and Mediterranean forest species: Direct effects and ecological consequences
E. Paoletti

Mediterranean plant adaptations against water stress and excess light may also afford protection against UV-B.

380 Possible impacts of changes in UV-B radiation on North American trees and forests
J.H. Sullivan

Projected increases in UV-B are not likely to have a direct impact on trees, but indirect responses could impact plant–herbivore interactions and nutrient cycling in forests.

390 UV-B radiation and acclimation in timberline plants
M. Turunen, K. Latola

More long-term field research is needed to assess the interaction of climate, soil and UV-B on timberline plants.

404 Growth and defense in deciduous trees and shrubs under UV-B
R. Julkunen-Tiitto, H. Häggman, P.J. Aphalo, A. Lavola, R. Tegelberg, T. Veteli

The growth performance of deciduous trees and shrubs under UV-B irradiation is constrained by multilevel interactions with many abiotic and biotic factors.

415 The effects of UVB radiation on temperate southern hemisphere forests
K.G. Ryan, J.E. Hunt

Solar UVB has a variable but generally modest effect on the morphology and growth of forest trees, but may modify forest community composition.

428 Depletion of stratospheric ozone over the Antarctic and Arctic: Responses of plants of polar terrestrial ecosystems to enhanced UV-B, an overview
J. Rozema, P. Boelen, P. Blokker

Polar plant responses to UV-B may be different in the Arctic than Antarctic regions.

443 Establishing a cause and effect relationship for ambient ozone exposure and tree growth in the forest: Progress and an experimental approach
W.J. Manning

Progress is reviewed and an experimental approach is proposed to demonstrate a cause and effect relationship for ambient ozone and forest tree growth.

455 Diagnosis of abiotic and biotic stress factors using the visible symptoms in foliage
P. Vollenweider, M.S. Günthardt-Goerg

The morphology and distribution of visible stress symptoms in tree foliage provides diagnostic tools to identify plant defense responses and differentiate stress from natural senescence symptoms.

466 Ozone-induced changes in the chloroplast structure of conifer needles, and their use in ozone diagnostics
M. Kivimäenpää, G. Selldén, S. Sutinen

Ozone-induced alterations in chloroplast structure of conifer needles are reviewed, and recommendations for field monitoring given.

476 Tree age dependence and within-canopy variation of leaf gas exchange and antioxidative defence in *Fagus sylvatica* under experimental free-air ozone exposure
K. Herbinger, C. Then, M. Löw, K. Haberer, M. Alexous, N. Koch, K. Remele, C. Heerdt, D. Grill, H. Rennenberg, K.-H. Häberle, R. Matyssek, M. Tausz, G. Wieser

Ozone effects on leaf gas exchange and antioxidative systems of beech across tree age and canopy level were investigated in a free air exposure system.

483 Does living in elevated CO_2 ameliorate tree response to ozone? A review on stomatal responses
E. Paoletti, N.E. Grulke

Understanding stomatal behavior under concurrent exposure to elevated CO_2 and O_3 is essential for predictive modeling of tree water balance.

494 Comparison of ozone uptake and sensitivity between a phytotron study with young beech and a field experiment with adult beech (*Fagus sylvatica*)
A.J. Nunn, A.R. Kozovits, I.M. Reiter, C. Heerdt, M. Leuchner, C. Lütz, X. Liu, M. Löw, J.B. Winkler, T.E.E. Grams, K.-H. Häberle, H. Werner, P. Fabian, H. Rennenberg, R. Matyssek

Juvenile beech trees in phytotrons are more sensitive to ozone than adult forest trees due to lower defence capacity and growth conditions.

507 Effects of simultaneous ozone exposure and nitrogen loads on carbohydrate concentrations, biomass, and growth of young spruce trees (*Picea abies*)
V.F.D. Thomas, S. Braun, W. Flückiger

Simultaneous ozone fumigation and nitrogen fertilization have no synergistic impacts on carbohydrate concentrations, biomass, or growth of Picea abies saplings.

517 Compensation processes of Aleppo pine (*Pinus halepensis* Mill.) to ozone exposure and drought stress
R. Inclán, B.S. Gimeno, P. Dizengremel, M. Sanchez

Ozone and drought stress induce the activation of similar processes related to C and N metabolism.

525 Canopy profiles of photosynthetic parameters under elevated CO_2 and N fertilization in a poplar plantation
C. Calfapietra, I. Tulva, E. Eensalu, M. Perez, P. De Angelis, G. Scarascia-Mugnozza, O. Kull

Photosynthetic acclimation occurred only marginally.

536 California black oak response to nitrogen amendment at a high O_3, nitrogen-saturated site
N.E. Grulke, W. Dobrowolski, P. Mingus, M.E. Fenn

Despite ecosystem-level N saturation, lower foliar respiration and significant photosynthetic gains under low light conditions resulted in greater wood production in black oak.

546 Air pollution, precipitation chemistry and forest health in the Retezat Mountains, Southern Carpathians, Romania
A. Bytnerowicz, O. Badea, F. Popescu, R. Musselman, M. Tanase, I. Barbu, W. Frączek, N. Gembasu, A. Surdu, F. Danescu, D. Postelnicu, R. Cenusa, C. Vasile

Retezat Mountains are characterized by relatively clean air, acidic precipitation, and healthy, well growing forests.

568 Modelling future soil chemistry at a highly polluted forest site at Istebna in Southern Poland using the "SAFE" model
S. Małek, L. Martinson, H. Sverdrup

Soil recovery depends on future emissions, especially on base cation deposition.

574 Fine roots in stands of *Fagus sylvatica* and *Picea abies* along a gradient of soil acidification
S. Braun, L. Cantaluppi, W. Flückiger

Fine root length of Fagus sylvatica and fine root depth in stands of Fagus sylvatica and/or Picea abies were impaired in soils with low base saturation.

580 Impacts of greenhouse gases on epicuticular waxes of *Populus tremuloides* Michx.: Results from an open-air exposure and a natural O_3 gradient
B. Maňkovská, K.E. Percy, D.F. Karnosky

Structure of epicuticular waxes indicated phytotoxic effects of greenhouse gases on Populus tremuloides Michx.

587 Meteorological factors and air pollution in Lithuanian forests: Possible effects on tree condition
R. Ozolincius, V. Stakenas, B. Serafinaviciute

Strongly damaged and relatively healthy trees respond differently to the same environmental factors.

596 The influence of climate and soil properties on calcium nutrition and vitality of silver fir (*Abies alba* Mill.)
N. Potočić, T. Ćosić, I. Pilaš

Dry years, especially in combination with calcium-poor soils, have a strong negative effect on the Ca status of silver fir trees.

603 Changes in the concentrations of phenolics and photosynthates in Scots pine (*Pinus sylvestris* L.) seedlings exposed to nickel and copper
M. Roitto, P. Rautio, R. Julkunen-Tiitto, E. Kukkola, S. Huttunen

Excess nickel in soil interferes with carbohydrate metabolism and induces an increase in concentration of condensed tannins in Scots pine needles.

610 Long-term evaluation of the needle surface wax condition of *Pinus sylvestris* around different industries in Lithuania
E. Kupcinskiene, S. Huttunen

Tubular wax on the pine needle surface reflects changes/differences in industrial emissions.